



DATE (dd/MMM/yy): \_\_\_\_\_ / \_\_\_\_\_ / \_\_\_\_\_



A. Type of Surgery	<input type="radio"/> Cardiac On Pump	<input type="radio"/> Cardiac Off Pump	<input type="radio"/> C-Section	<input type="radio"/> General Surgery	<input type="radio"/> Gynecology	<input type="radio"/> Ophthalmology	<input type="radio"/> Orthopedic	<input type="radio"/> Thoracic	<input type="radio"/> Vascular	<input type="radio"/> Other
B. Surgical Class	<input type="radio"/> Clean (I)	<input type="radio"/> Clean-Contaminated (II)	<input type="radio"/> Contaminated (III)	<input type="radio"/> Dirty (IV)	<input type="radio"/> Not Recorded					
C. Pre-Op shower or bath with soap or antiseptic agent	<input type="radio"/> Soap	<input type="radio"/> Antiseptic	<input type="radio"/> Shower or bath not required	<input type="radio"/> No shower or bath	<input type="radio"/> Not Recorded					
D. Solution used for intra-operative intact skin cleansing	<input type="radio"/> 2% Chlorhexidine in 70% Alcohol	<input type="radio"/> Povidone-Iodine with Alcohol	<input type="radio"/> Povidone-Iodine for Head & Neck	<input type="radio"/> Contraindicated	<input type="radio"/> Not Recorded					
	<input type="radio"/> Chlorhexidine	<input type="radio"/> Povidone-Iodine	<input type="radio"/> Other	<input type="radio"/> Not Applicable						
E. Prophylactic Abx administration	<input type="radio"/> Within 60 minutes before incision	<input type="radio"/> Within 120 minutes before incision for Vancomycin or Fluroquinolones	<input type="radio"/> None of the above	<input type="radio"/> No Abx given	<input type="radio"/> Antibiotics not indicated	<input type="radio"/> Not Recorded				
			<input type="radio"/> Not Recorded							
F. Dose of Cefazolin used as prophylactic Abx (Adults only)	<input type="radio"/> Weight based dose for pediatric pt	<input type="radio"/> < 1g for any adult pt	<input type="radio"/> Other Abx used							
	<input type="radio"/> 1g <80kg	<input type="radio"/> 1g >=80kg	<input type="radio"/> 2g <80kg	<input type="radio"/> 2g 80-120kg	<input type="radio"/> 2g >120kg					
	<input type="radio"/> 3g <120kg	<input type="radio"/> 3g >=120kg	<input type="radio"/> >3g	<input type="radio"/> No Abx given	<input type="radio"/> Not Recorded					
G. Appropriate prophylactic Abx redosing according to guidelines	<input type="radio"/> No prophylactic antibiotic given	<input type="radio"/> Yes	<input type="radio"/> No	<input type="radio"/> Redosing not required	<input type="radio"/> Not Recorded					
H. Discontinuation of prophylactic Abx	<input type="radio"/> Abx not received after end of surgery	<input type="radio"/> Abx discont'd within 24 hrs of end of surgery	<input type="radio"/> Abx discont'd more than 24 hrs after end of surgery	<input type="radio"/> No Abx given	<input type="radio"/> Not Recorded					
I. Hair removal method	<input type="radio"/> Hair removal not req'd	<input type="radio"/> Clippers	<input type="radio"/> Depilatory	<input type="radio"/> Razor	<input type="radio"/> Removal done at home	<input type="radio"/> Not Recorded				
J. Glucose was below 11.1 mmol/L on each of POD 0, 1, & 2	<input type="radio"/> Not at risk	<input type="radio"/> Yes	<input type="radio"/> No	<input type="radio"/> Glucose not done	<input type="radio"/> Not Recorded					
K. Temp at end of surgery or on arrival in PACU was within range of 36.0-38.0 C	<input type="radio"/> Yes	<input type="radio"/> No	<input type="radio"/> Induced hypothermia	<input type="radio"/> Not recorded						

Question	C.	D.	E.	F.	G.	H.	I.	J.	K.
<b>Best Practice (Score 0 or 1)</b>									

Refer to the detailed instructions for directions for completing the data collection form and calculating the results. Instructions are available in the measurement package from Safer Healthcare Now! (metrics@saferhealthcarenow.ca)

